

AD-A125 106

SYNTHESIS AND CHARACTERIZATION OF ORGANOALUMINUM  
COMPOUNDS CONTAINING THE (U) STATE UNIV OF NEW YORK AT  
BUFFALO DEPT OF CHEMISTRY O T BEACHLEY ET AL  
08 FEB 83 TR-10 N00014-78-C-0562

1/1

UNCLASSIFIED


F/G 7/3

NL

END

FILMED

100%



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD A125106

OFFICE OF NAVAL RESEARCH  
Contract N-00014-78-C-0562  
Task No. NR 053-686  
TECHNICAL REPORT NO. 10

Synthesis and Characterization of Organoaluminum Compounds  
Containing the Trimethylsilylmethyl Substituent  
 $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$ ,  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  and  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$  and  
a Reinvestigation of the Chemistry of  $\text{Me}_2\text{AlPPh}_2$  and  $\text{Et}_2\text{AlPPh}_2$ .

by

O. T. Beachley, Jr. and Claire Tessier-Youngs


Prepared for Publication

in

Organometallics

State University of New York at Buffalo  
Department of Chemistry  
Buffalo, New York 14214

8, February 1983



Reproduction in whole or in part is permitted for any purpose of the  
United States Government

\*This document has been approved for public release and sale; its  
distribution is unlimited

83 02 017 013

DTIC FILE COPY

(Contribution from the Department of Chemistry,  
State University of New York at Buffalo, Buffalo, New York 14214)


Synthesis and Characterization of Organoaluminum Compounds  
Containing the Trimethylsilylmethyl Substituent,  
 $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$ ,  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  and  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$  and  
a Reinvestigation of the Chemistry of  $\text{Me}_2\text{AlPPh}_2$  and  $\text{Et}_2\text{AlPPh}_2$ .

by

O. T. Beachley, Jr.\* and Claire Tessier-Youngs

Abstract

The new amphoteric ligand,  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$ , has been prepared from the reactions of either  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$  or  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  with  $\text{PPh}_2\text{H}$  or  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  with  $\text{KPPH}_2$  and fully characterized by analysis, cryoscopic molecular weight measurements, IR and  $^1\text{H}$  NMR data. This aluminum-phosphide is unique as it exists as a monomer-dimer equilibrium mixture in benzene solution. The syntheses and characterization of the new compounds  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  and  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  are also described. Since the chemistry of aluminum-phosphides as amphoteric ligands is of interest, the behavior of  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$ ,  $(\text{Me}_2\text{AlPPh}_2)_2$  and  $(\text{Et}_2\text{AlPPh}_2)_2$  toward common solvents,  $\text{Et}_2\text{O}$ , THF and  $\text{CH}_3\text{CN}$ , was investigated. The aluminum phosphides readily cleave THF and reduce the triple bond of  $\text{CH}_3\text{CN}$  at room temperature. However, neither THF nor  $\text{Et}_2\text{O}$  form isolable adducts with aluminum phosphides. The unusual melting point behavior of aluminum phosphides is also discussed.



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Dissemination	
Availability Codes	
Distribution Statement	
Final	
A	

## Introduction

The synthesis and characterization of  $\text{Cr}(\text{CO})_5[\text{PPh}_2\text{Al}(\text{CH}_2\text{SiMe}_3)_2\cdot\text{NMe}_3]^1$  has been described recently. This compound serves as a model for compounds in which a diphenylphosphide group bridges a transition metal and a main-group element moiety. Alternatively, this new compound may be considered to be a transition metal derivative of an organoaluminum phosphide, an amphoteric ligand. As part of this research effort, the chemistry of amphoteric ligands is of interest and consequently we have extended, reinvestigated and in some cases reassessed the chemistry of several organoaluminum diphenylphosphides.

Organoaluminum diphenylphosphides with methyl,<sup>2</sup> ethyl<sup>3</sup> and aryl<sup>4</sup> substituents have been prepared from triorganoaluminum compounds and  $\text{PPh}_2\text{H}$  by elimination reactions. The compounds  $\text{R}_2\text{AlPPh}_2$  exist as dimers in benzene solution and are believed to have four-membered aluminum-phosphorus rings in analogy with similar compounds containing other Group 3 and Group 5 atoms.<sup>2-4</sup> The four-membered rings reportedly can be disrupted by reaction with various Lewis bases to form adducts of formula  $\text{Ph}_2\text{PAIR}_2\cdot\text{Base}$ .<sup>2-4</sup> However, no NMR or x-ray structural data have been reported to support the proposed structures of dimers or of their adducts. The lack of such data prompted us to undertake a reinvestigation of the properties and reaction chemistry of  $\text{Me}_2\text{AlPPh}_2$  and  $\text{Et}_2\text{AlPPh}_2$ . The results of our initial attempts to synthesize  $\text{M}(\text{CO})_5\text{PPh}_2\text{AlR}_2$  compounds defined the need for a more soluble organoaluminum diphenylphosphide. Our previous experience<sup>5,6</sup> had indicated that the trimethylsilylmethyl ligand enhanced the solubility of its compounds in hydrocarbon solvents. Thus, we prepared  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$  from  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$ ,  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  and  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  in order to define the best and most useful route. Our new convenient synthesis of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$  has been reported.<sup>7</sup>

From this parent compound, the preparations of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  and  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  have been accomplished. In this paper we also report the properties of the adducts  $\text{R}_3\text{AlPPh}_2\text{H}$  ( $\text{R} = \text{Me}, \text{Et}, \text{CH}_2\text{SiMe}_3$ ) and the chemistry of organoaluminum diphenylphosphides towards ethers (THF and  $\text{Et}_2\text{O}$ ) and  $\text{CH}_3\text{CN}$ .

### Experimental

Handling procedures, solvent purification and spectra were as previously described.<sup>1</sup> All molecular weights were measured cryoscopically in benzene solution.<sup>1</sup> Moles of hydrolyzable alkyl groups or hydrogens bound to aluminum were determined by measuring the gas (PVT) evolved upon acid hydrolysis from a weighed sample of compound<sup>1</sup> (mol). Attempts to analyze for aluminum via an EDTA titration<sup>2</sup> in the presence of phosphorus were unsuccessful. Immediately prior to its use  $\text{AlMe}_3$  was vacuum distilled.  $\text{AlEt}_3$  was obtained from a commercial 25% solution in hexane by removal of hexane by vacuum distillation. The compound  $\text{AlMe}_2\text{H}$  was obtained from the reaction of  $\text{AlMe}_3$  and  $\text{LiAlH}_4$ <sup>8</sup> and was purified by vacuum distillation. The reagents  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$ <sup>7</sup> and  $\text{PPh}_2\text{H}$ <sup>9</sup> were synthesized by literature procedures. The liquids,  $\text{AlEt}_3$  (0.835 g/mL)<sup>10</sup> and  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$  (0.802 g/mL),<sup>7</sup> were measured by volume using graduated gas-tight syringes in the dry-box. The liquid  $\text{PPh}_2\text{H}$  (1.07 g/mL)<sup>11</sup> was similarly handled using Schlenk techniques. <sup>1</sup>H NMR spectra are reported in  $\delta$  units (ppm) and are referenced to TMS as 0.00 and  $\text{C}_6\text{H}_6$  as 7.27. Assignments are based on integration of spectra, where possible.

#### Synthesis of $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$

The compound  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  was synthesized by an exchange reaction between one mol  $\text{AlBr}_3$  and two mol  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$ . In the dry box, 1.75 mL (1.40 g, 4.87 mmol)  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$  and 0.650 g (2.43 mmol)  $\text{AlBr}_3$  were combined in

a 10 mL flask. Heat was evolved and a colorless liquid was produced. The flask was attached to a micro-scale distillation apparatus and  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  (1.926 g, 94%) distilled as a colorless liquid at 60-62°/0.01 mm. Density: 1.06 g/mL at 22°. Anal. Calc.: Al, 9.59; Br, 28.41;  $\text{Me}_4\text{Si}$ , 2.00 mol/mol. Found: Al, 9.45; Br, 27.73;  $\text{Me}_4\text{Si}$ , 1.98 mol/mol.  $^1\text{H}$  NMR (benzene,  $\delta$ ): 0.27 (s, 9,  $\text{CH}_3$ ), -0.16 (s, 2,  $\text{CH}_2$ ). IR (Nujol,  $\text{cm}^{-1}$ ): 1259(s), 975(s,b), 946(m), 858(vs,b), 763(s), 734(s), 668(m), 636(vs), 577(s), 315(w). Cryoscopic molecular weight. Formula weight  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$ : 281.27. Molality, observed mol. wt., association: 0.1257, 644, 2.29; 0.0497, 405, 1.44; 0.0401, 335, 1.19; 0.0336, 295, 1.05. Solubility: soluble in aromatic, halogenated and aliphatic hydrocarbons and ethers.  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  readily reacts with absorbed water on glass. In order to obtain satisfactory analyses for this compound, the reaction flask had to be oven dried (>100°C) prior to being brought into the dry box and the flask containing the sample had to be evacuated at -196° to avoid loss of volatile  $\text{Me}_4\text{Si}$  and  $\text{HBr}$  which formed even with these precautions.

#### Synthesis of $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$

The reagents  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$  (2.5 mL, 2.0 g, 6.9 mmol) and 0.3 g (7.9 mmol)  $\text{LiAlH}_4$  (recrystallized from  $\text{Et}_2\text{O}$ ) were heated at 90° for 18 h under vacuum. The reaction mixture solidified on cooling to room temperature. The product  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  (1.29 g, 92%) sublimed at 40-70° under high vacuum as a crystalline solid. Mp: 35-36°. Alternatively,  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  was isolated from the reaction mixture by extracting with hexane, concentrating the solution and cooling to 0° or -20°. Mp: 38-40°. Anal. Calc.: Al, 13.33;  $\text{Me}_4\text{Si}$ , 2.00 mol/mol;  $\text{H}_2$ , 1.00 mol/mol. Found: Al, 11.36;  $\text{Me}_4\text{Si}$ , 2.04 mol/mol;  $\text{H}_2$  0.99 mol/mol.  $^1\text{H}$  NMR

(benzene,  $\delta$ ):<sup>1</sup> 0.33 (s, 9, CH<sub>3</sub>), -0.25 (s, 2, CH<sub>2</sub>). IR (Nujol mull, cm<sup>-1</sup>): 1765 (vs, vb, Al-H), 1257 (vs), 970 (vs, b), 854 (vs), 838 (vs), 765 (vs), 733 (s), 702 (m), 661 (w), 575 (m). Cryoscopic molecular weight. Formula weight Al(CH<sub>2</sub>SiMe<sub>3</sub>)<sub>2</sub>H: 202.35. Molality, obs. mol. wt., association: 0.1144, 607, 3.00; 0.144, 648, 3.20; 0.1702, 609, 3.01. Solubility: soluble in aromatic and aliphatic hydrocarbons. The analysis suggests a 5-9% impurity of Al(CH<sub>2</sub>SiMe<sub>3</sub>)<sub>3</sub> which could not be removed due to the similarities of its solubility and volatility with Al(CH<sub>2</sub>SiMe<sub>3</sub>)<sub>2</sub>H.

#### Synthesis of R<sub>3</sub>AlPPh<sub>2</sub>H (R = Me, Et, CH<sub>2</sub>SiMe<sub>3</sub>)

Adducts of formula R<sub>3</sub>Al·PPh<sub>2</sub>H (R = Me, Et or Me<sub>3</sub>SiCH<sub>2</sub>) were obtained by combining equimolar quantities of the trialkylaluminum compound and Ph<sub>2</sub>PH (see preparations of R<sub>2</sub>AlPPh<sub>2</sub>). All three adducts were colorless liquids at room temperature. <sup>1</sup>H NMR Me<sub>3</sub>Al·PPh<sub>2</sub>H (d<sup>8</sup>-toluene,  $\delta$ ): 7.46 (m, Ph), 7.20 (m, Ph), 5.53 (d, J = 299 Hz, 1, PH), -0.08 (s, 9, CH<sub>3</sub>). <sup>1</sup>H NMR Et<sub>3</sub>Al·PPh<sub>2</sub>H (d<sup>6</sup>-benzene,  $\delta$ ): 7.45 (m, Ph), 7.18 (m, Ph), 5.55 (d, J = 295 Hz, 1, PH), 1.51 (t, J = 8 Hz, 9, CH<sub>3</sub>), 0.53 (q, J = 8 Hz, 6, CH<sub>2</sub>). <sup>1</sup>H NMR (Me<sub>3</sub>SiCH<sub>2</sub>)<sub>3</sub>Al·PPh<sub>2</sub>H (d<sup>6</sup>-benzene,  $\delta$ ): 7.55 (m, Ph), 7.24 (m, Ph), 5.62 (d, J = 300 Hz, 1, PH), 0.30 (s, 27, CH<sub>3</sub>), -0.40 (s, 6, CH<sub>2</sub>).

#### Synthesis of (Me<sub>3</sub>SiCH<sub>2</sub>)<sub>2</sub>AlPPh<sub>2</sub>

##### a) From Ph<sub>2</sub>PH and Al(CH<sub>2</sub>SiMe<sub>3</sub>)<sub>3</sub>

As in the preparation of Et<sub>2</sub>AlPPh<sub>2</sub>, a break-seal tube containing 1.00 mL (1.07 g, 5.74 mmol) Ph<sub>2</sub>PH and 2.09 mL (1.68 g, 5.82 mmol) Al(CH<sub>2</sub>SiMe<sub>3</sub>)<sub>3</sub> was prepared. The tube was evacuated, sealed at -196° and then heated at 160-180° for about 18 h. The progress of the



reaction was followed by periodically cooling the tube to room temperature. At room temperature,  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$  is a solid and can be distinguished from the liquid adduct. The tube was opened under vacuum. Tetramethylsilane (5.45 mmol, 94.9%) and a trace of a noncondensable gas were evolved during the reaction. The solid was transferred from the tube to a flask by repeated extractions with one 5 mL portion of hexane and then washed once with the same hexane at  $-20^\circ$  to remove unreacted adduct. The compound  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$  (1.63 g, 73.4%) was obtained as a white solid. Mp:  $120-121^\circ$  a glass forms which melts at  $136-139^\circ$ ; cooling this sample to room temperature gives crystals which when slowly heated melt at  $141-143^\circ$ .  $^1\text{H}$  NMR (benzene,  $\delta$ ): 0.20 (s, 4.5,  $\text{CH}_3$ ), 0.06 (q,  $J = 2.7$  Hz, 1,  $\text{CH}_2$ ). (See Results and Discussion for additional description of spectrum.) IR (Nujol mull,  $\text{cm}^{-1}$ ): 1584 (m), 1263 (m), 1248 (s), 1097 (m,b), 1034 (m), 971 (s), 956 (m), 862 (vs), 832 (vs), 749 (vs), 733 (s,sh), 704 (m), 696 (m,sh), 652 (m), 636 (w), 567 (m), 470 (w), 451 (w), 430 (w), 398 (w), 313 (w). Cryoscopic molecular weight. Formula weight  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$ : 386.54. Molality, obs. mol. wt., association: 0.0549, 665, 1.72; 0.0446, 638, 1.65; 0.0210, 545, 1.41.

b) From  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  and  $\text{Ph}_2\text{PH}$

A sample of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  (0.660 g, 3.26 mmol) was placed in a tube equipped with a Teflon valve. Toluene (5 mL) was distilled into the tube and then, under argon flush, the tube was attached to a two-neck flask containing 0.56 mL (0.60 g, 3.2 mmol)  $\text{Ph}_2\text{PH}$ . The flask was evacuated and the alane solution was added to the stirred phosphine. Evolution of a noncondensable gas took place very slowly at room

temperature in toluene solution. The toluene was removed by vacuum distillation and the reaction flask was heated with an oil bath. Gas evolution was rapid at 65° and a white solid formed. The flask was heated at 70-80° for 2 h to drive the reaction to completion. Hydrogen was evolved in greater than 91% yield. Unreacted adduct was removed by washing the solid with 5 mL hexane at -20°. Mp: 146-147°. Anal. Calc.:  $\text{Me}_4\text{Si}$ , 2.00 mol/mol. Found:  $\text{Me}_4\text{Si}$ , 2.02 mol/mol. Infrared and  $^1\text{H}$  NMR spectra were identical to those obtained from the previous reaction.

c) From  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  and  $\text{KPPH}_2$

Potassium hydride (0.102 g, 2.54 mmol) was slowly added to a stirred solution of 0.64 mL (0.69 g, 3.7 mmol)  $\text{Ph}_2\text{PH}$  in 10 mL diethyl ether. An orange solid,  $\text{KPPH}_2$ , was precipitated and hydrogen was evolved in 95% yield in less than 12 hr. A solution of 0.97 mL (1.0 g, 3.7 mmol)  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  in 3 mL diethyl ether, contained in a built-in addition tube of the reaction vessel, was added to the stirred  $\text{KPPH}_2$  suspension. The orange solid disappeared immediately and a white precipitate was formed. After stirring for one-half hour, the solution was filtered from the KBr using a medium frit. Diethyl ether was removed by vacuum distillation and the remaining white solid was recrystallized from pentane. The compound  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPH}_2$  was obtained in 71% yield. Mp: glass forms at 115° which melts completely by 138°. The  $^1\text{H}$  NMR spectrum of the product in  $\text{d}^6$ -benzene indicated  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPH}_2$  and the presence of a small impurity of  $\text{Br}(\text{Me}_3\text{SiCH}_2)_2\text{AlPPH}_2\text{H}$ .

## Syntheses of $\text{Me}_2\text{AlPPh}_2$

### a) From $\text{PPh}_2\text{H}$ and $\text{AlMe}_3$

The compound  $\text{AlMe}_3$  (1.0117 g, 14.03 mmol) was distilled into a trap on the vacuum line. Then  $\text{Ph}_2\text{PH}$  (2.34 mL, 2.50 g, 13.4 mmol) was syringed into an argon filled break-seal tube. The preweighed  $\text{AlMe}_3$  was vacuum distilled into the tube and the tube was sealed at  $-196^\circ$ . On warming to room temperature, an exothermic reaction took place. The clear liquid adduct was heated for 16 h at  $160^\circ$ , producing a white solid and 13.4 mmol methane gas. The tube was opened under vacuum using a stainless steel breaker. The contents of the tube were transferred into a flask by repeated extractions with toluene. The resultant white solid was washed once with hexane (5 mL) yielding  $\text{Me}_2\text{AlPPh}_2$  (2.812 g, 86%) as a white powder. Mp:  $200-206^\circ$  a glass forms,  $211-222^\circ$  the glass melts (lit. mp:  $230^\circ$ , sublimed).<sup>2</sup> Anal. Calc.: MeH, 2.00 mol/mol. Found: MeH, 1.99 mol/mol.  $^1\text{H}$  NMR ( $\text{d}^8$ -toluene,  $\delta$ ): 7.50 (m, Ph), 7.16 (m, Ph), 0.08 (q,  $J = 2$  Hz,  $\text{AlCH}_3$ ). (See Results and Discussion for additional description of spectrum.) IR (Nujol mull,  $\text{cm}^{-1}$ ): 3081 (w), 3050 (w), 1587 (w), 1437 (s), 1264 (w), 1183 (m), 1042 (s,b), 1021 (s), 1005 (s), 950 (w), 926 (w), 804 (w), 757 (m), 730 (s), 685 (s,b), 554 (m), 487 (m), 423 (m), 355 (w), 332 (w). Solubility: moderately soluble in benzene or toluene, insoluble in hexane and diethyl ether. The compound  $\text{Me}_2\text{AlPPh}_2$  reacts with halogenated hydrocarbons, THF and  $\text{CH}_3\text{CN}$  (vide infra), solvents in which it has only slight solubility. The attempted sublimation of a 1.79 g sample at  $160^\circ$  for 12 h under high vacuum produced 0.0347 g (2%) sublimed  $\text{Me}_2\text{AlPPh}_2$  (mp:  $228^\circ$ ).

b) From  $\text{Ph}_2\text{PH}$  and  $\text{AlMe}_3$  in toluene.

As in the previous preparation, a break-seal tube containing 0.8315 g (11.53 mmol)  $\text{AlMe}_3$  and 1.54 mL (1.65 g, 8.88 mmol)  $\text{Ph}_2\text{PH}$  was prepared. Toluene (10 mL) was vacuum distilled into the tube and the tube was sealed. The reaction mixture was heated at  $140^\circ$  for 18 h. Large crystals formed as the reaction vessel slowly cooled to room temperature. The tube was opened under vacuum. Methane was generated by the reaction. The crystals were washed out of the tube with toluene, recrystallized from this solvent and subjected to high vacuum for several hours. During this latter process the unreacted  $\text{AlMe}_3$  was removed. The compound  $\text{Me}_2\text{AlPPh}_2$  (1.40 g, 62%) was obtained as a white solid which melted at  $228^\circ$ . Identical  $^1\text{H}$  NMR and infrared spectra were obtained as for the compound described in the previous preparation.

c) From  $\text{Ph}_2\text{PH}$  and  $\text{AlMe}_2\text{H}$ .

The reagent  $\text{AlMe}_2\text{H}$  (0.2789 g, 4.804 mmol) was vacuum distilled into a tube equipped with a Teflon stopcock. Under argon flush, the tube was attached to a 2-neck flask containing 0.83 mL (0.88 g, 4.8 mmol)  $\text{Ph}_2\text{PH}$ . The flask was evacuated and  $\text{AlMe}_2\text{H}$  was transferred to the flask by distillation. The reagents were allowed to slowly warm to room temperature. Vigorous bubbling took place and a white solid and a noncondensable gas formed. After 2 h at room temperature, the reaction flask was heated at  $90^\circ$  for 1 h to drive the reaction to completion. Using estimated volumes, the quantity of hydrogen evolved was nearly quantitative. A sublimation of this solid was attempted but after several hours at  $160^\circ$  only a trace of a liquid, identified as  $\text{Ph}_2\text{PH}$ , had distilled to the cold finger. The remaining solid was then recrystallized from toluene, yielding 0.235 g

(20%)  $\text{Me}_2\text{AlPPh}_2$ . Mp: 210-212° glass forms, which is completely melted by 225°. Identical infrared and  $^1\text{H}$  NMR spectra were obtained as for the compounds described in the previous two preparations. A second crop of a slightly sticky solid (0.20 g) melting at 205-210°, was shown to contain  $\text{Ph}_2\text{PH}$  by its infrared spectra.

#### Synthesis of $\text{Et}_2\text{AlPPh}_2$

The reagent  $\text{Ph}_2\text{PH}$  (2.25 mL, 2.41 g, 2.9 mmol) was syringed into an argon filled break-seal tube. In the dry-box, 1.79 mL (1.49 g, 13.0 mmol)  $\text{AlEt}_3$  was added to the tube. Heat was evolved due to adduct formation. Toluene (10 mL) was then vacuum distilled into the tube. The tube was sealed at -196° and then heated at 90° for 16 h. The tube was opened under vacuum and the ethane evolved (11.4 mmol, 88.4%) was measured. The toluene solution was filtered into a flask under vacuum. Solvent was removed by vacuum distillation leaving a sticky white solid. The solid was recrystallized from 6 mL 1:1 toluene/hexane yielding  $\text{Et}_2\text{AlPPh}_2$  (2.374 g, 68%) as a white powder. Mp: 123-126° glass forms, 141-143° glass melts (lit. mp: 124-126°, from benzene).<sup>3</sup> In other preparations, the sticky white solid was washed with hexane, leaving a solid which melted at 140-143°. Identical spectral and analytical data were obtained for both products.

Anal. Calc.: EtH, 2.00 mol/mol. Found: EtH, 2.02 mol/mol.  $^1\text{H}$  NMR ( $d^8$ -toluene,  $\delta$ ): 7.51 (m, Ph), 7.20 (m, Ph), 1.31 (t,  $J = 7$  Hz, 1.5,  $\text{CH}_3$ ), 0.80 (q,  $J = 7$  Hz, 1,  $\text{CH}_2$ ). (See Results and Discussion for additional description of spectrum.)  $^{31}\text{P}$  NMR ( $d^8$ -toluene, ppm from 85%  $\text{H}_3\text{PO}_4$ ): -42.30 (s). IR (Nujol mull,  $\text{cm}^{-1}$ ): 3052 (w), 1594 (w), 1441 (s), 1267 (w), 1236 (w), 1195 (m), 1137 (vs), 1051 (m,b), 1018 (m), 989 (m), 955 (m), 920 (w), 809 (w), 759 (m), 736 (s), 700 (s), 543 (m), 504 (m), 477 (m), 424 (w).

400 (w), 379 (vw), 284 (vw). Solubility: soluble in aromatic hydrocarbons and diethyl ether, slightly soluble in hexane, soluble in and reacts with halogenated hydrocarbons, THF and  $\text{CH}_3\text{CN}$  (vide infra).

#### Attempted Syntheses of $\text{Et}_2\text{AlPPh}_2 \cdot \text{OEt}_2$

A 0.3180 g (1.177 mmol) sample of  $\text{Et}_2\text{AlPPh}_2$  was placed in a tube equipped with a Teflon valve and a magnetic stirring bar. The tube was evacuated and weighed. Diethyl ether (2 mL) was vacuum distilled into the tube. The solution was stirred until all the  $\text{Et}_2\text{AlPPh}_2$  dissolved and then cooled to  $-78^\circ$  for 48 h.  $\text{Et}_2\text{O}$  was removed by vacuum distillation and the tube was reweighed. No diethyl ether was retained by the sample. This observation was confirmed by the  $^1\text{H}$  NMR spectrum which identified the presence of only  $\text{Et}_2\text{AlPPh}_2$ . The sample melted at  $143\text{--}146^\circ$  (lit. mp  $\text{Et}_2\text{AlPPh}_2 \cdot \text{OEt}_2$ :  $148^\circ$ ).<sup>3</sup>

Another sample of  $\text{Et}_2\text{AlPPh}_2$  was recrystallized in  $\text{Et}_2\text{O}$ . The crystals were subjected to high vacuum at  $-78^\circ$  until they appeared dry. A  $^1\text{H}$  NMR spectrum of these crystals showed that no ether was retained. The crystals melted at  $143\text{--}145^\circ$ .

#### Reaction of $\text{Et}_2\text{AlPPh}_2$ with THF

A sample of  $\text{Et}_2\text{AlPPh}_2$  and THF was stirred for 18 h in a flask equipped with a side-arm NMR tube. Solvent was removed by vacuum distillation and some of the resultant sticky white solid was washed into the NMR tube with benzene. The tube was sealed at  $-196^\circ$  and the  $^1\text{H}$  NMR spectrum was recorded.  $^1\text{H}$  NMR (benzene,  $\delta$ ): 3.55 (t,  $J = 6$  Hz,  $\text{OCH}_2$ ), 2.02 (t,  $J = 7$  Hz,  $\text{PCH}_2$ ), 1.44 (m, other  $\text{CH}_2$ ), 1.36 (t,  $J = 9$  Hz,  $\text{AlCH}_2\text{CH}_3$ ), 0.32 (m, b,  $\text{AlCH}_2$ ), 0.21 (q,  $J = 9$  Hz,  $\text{AlCH}_2$ ). The sticky material remaining in the reaction tube was then treated with dilute hydrochloric acid and extracted with diethyl ether. Solvent was removed from the extract, the clear oil was dried under

high vacuum for three days and its infrared spectrum was recorded.

$\text{Ph}_2\text{PC}_4\text{H}_8\text{OH}$  plus  $\text{PPh}_2\text{H}$ . IR (neat, bands above  $1400\text{ cm}^{-1}$ ): 3380 (vs, b, OH), 3058 (s), 2964 (s), 2932 (s), 2866 (s), 2332 (m, PH); 1969 (w), 1899 (w), 1821 (w), 1720 (m), 1592 (m), 1583 (s), 1490 (m), 1442 (s).

#### Reactions of Organoaluminum Phosphides with Acetonitrile


Samples of organoaluminum phosphides (about 0.1 g) were stirred in  $\text{CH}_3\text{CN}$  (3 mL) for 48 h. The solvent was removed by vacuum distillation. Sticky white solids, which were soluble in toluene, were obtained.

Infrared spectra of Nujol mulls were recorded. Bands assignable to carbon-nitrogen double or triple bonds are reported.  $\text{Me}_2\text{AlPPh}_2 + \text{CH}_3\text{CN}$ .

Mp:  $80^\circ$  red color,  $138\text{--}140^\circ$  sample melts to a red liquid. IR ( $\text{cm}^{-1}$ ):  $\nu_{\text{C=N}}$  1601 (s).  $\text{Et}_2\text{AlPPh}_2 + \text{CH}_3\text{CN}$ . Mp:  $125^\circ$  glass forms,  $137^\circ$  sample melts to a yellow liquid. IR ( $\text{cm}^{-1}$ ):  $\nu_{\text{C=N}}$  1613 (s).  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2 + \text{CH}_3\text{CN}$ . Mp:  $160\text{--}165^\circ$  to a yellow liquid. IR ( $\text{cm}^{-1}$ ):  $\nu_{\text{C=N}}$  2186 (m),  $\nu_{\text{C=N}}$  1601 (s).

#### Results and Discussion

Standard reactions have been used for the preparation of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  and  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  from the trialkylaluminum compound. The compound  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  was prepared in 94% yield by an exchange reaction between  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$  and  $\text{AlBr}_3$  in the absence of solvent (eq. 1).




Exchange is exothermic and apparently occurs at room temperature.

Formulation of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  as a discrete compound, rather than as a mixture of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$  and  $\text{AlBr}_3$  is supported by the fact that  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  has a two degree boiling range about ten degrees higher than  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$  under

high vacuum. The compound  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  exists as a mixture of monomeric and dimeric species in benzene solution in contrast to the dimeric association observed in hydrocarbon solutions of other organoaluminum halides,<sup>12</sup>  $\text{In}(\text{CH}_2\text{SiMe}_3)_2\text{Cl}$ <sup>6</sup> and  $\text{Ga}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$ .<sup>5</sup> The lower association of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$  in comparison to these compounds can be attributed to a combination of the large steric size of the trimethylsilylmethyl substituents and the smaller size of the aluminum atom. The analogous chloro-substituted compound,  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Cl}$ , was prepared from the reaction of one mole of  $\text{AlCl}_3$  with two moles of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$ .<sup>13</sup> The solution molecular weight of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Cl}$  was not reported but mass spectral data showed a fragmentation pattern consistent with a monomeric species in the gas phase.

The synthesis of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  has been achieved by the reaction of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$  with an excess of  $\text{LiAlH}_4$  at  $90^\circ$  (eq. 2). This type of




reaction has also been used to prepare  $\text{AlMe}_2\text{H}$ .<sup>8</sup> The compound  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  can be easily isolated in about 90% yield by sublimation or crystallization from hexane. However analysis of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  indicates the presence of a 5-9% impurity of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$ , which cannot be removed by repeated sublimations or crystallizations due to similarities of volatility and solubility properties of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  and  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$ . This situation was not improved by altering the reaction conditions. Under the preparative reaction conditions,  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$  apparently distills away from the  $\text{LiAlH}_4$  to cooler parts of the reaction vessel to avoid reaction. The reaction has also been attempted several times using hexane or toluene as solvent in the hope that the solvent would wash unreacted  $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$  back down onto



the  $\text{LiAlH}_4$ . Surprisingly, only very impure solids were isolated from these reactions. Use of longer reaction times or a larger excess of  $\text{LiAlH}_4$  were also ineffective in reducing the amount of the impurity.

The compound  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  is the only dialkylaluminum hydride that exists as a solid at room temperature. Other dialkylaluminum hydrides, with alkyl groups ranging in size from methyl to iso-butyl, exist as viscous oils or as liquids.<sup>12</sup> Cryoscopic molecular weight data indicate that  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  is trimeric in benzene solution. A puckered 6-membered ring composed of alternating dialkylaluminum groups and hydrogen atoms has been proposed for the structure of other trimeric  $\text{AlR}_2\text{H}$  compounds ( $\text{R} = \text{Me, et, iBu}$ ).<sup>12</sup> The infrared spectrum of  $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$  shows a very broad and intense band





centered at  $1765\text{ cm}^{-1}$  which can be attributed to the aluminum-hydrogen stretching frequency. Similarly broad and intense bands in the range of  $1785\text{-}1775\text{ cm}^{-1}$  have been observed in the infrared spectra of other dialkylaluminum hydrides.<sup>14</sup>

The main route by which the compounds  $\text{R}_2\text{AlPPh}_2$  ( $\text{R} = \text{Me, Et, CH}_2\text{SiMe}_3$ ) have been synthesized involves an elimination-condensation reaction between  $\text{AlR}_3$  and  $\text{Ph}_2\text{PH}$ . Formation of the intermediate adduct is an exothermic process giving a liquid product in all three cases. The adducts  $\text{R}_3\text{Al}\cdot\text{PPh}_2\text{H}$  have been examined by  $^1\text{H}$  NMR spectroscopy. Complexation of diphenylphosphine by aluminum is shown by the P-H  $^1\text{H}$  NMR signal, a doublet due to  $^{31}\text{P}\text{-}^1\text{H}$  coupling.<sup>15</sup> The chemical shift for the P-H of  $\text{Ph}_2\text{PH}$  ( $\delta$  5.2-5.1) depending on solvent

moves to lower field upon coordination to the trialkylaluminum compounds ( $\delta = 5.7-5.5$ ). The solvent dependency of the chemical shift makes it a less useful probe of complexation than the  $^{31}\text{P}-^1\text{H}$  coupling constant, which is not only virtually solvent independent but is also highly sensitive to the degree of substitution on the phosphorus atom.<sup>15</sup> In aromatic solvents, the  $^{31}\text{P}-^1\text{H}$  coupling constant of  $\text{PPh}_2\text{H}$  is about 220 Hz. This value is in the 180-230 Hz range indicative of three coordinate phosphorus.<sup>15</sup> The  $^1\text{H}$  NMR spectra of  $\text{R}_3\text{AlPPh}_2\text{H}$  in aromatic solvents show a substantial increase in the  $^{31}\text{P}-^1\text{H}$  coupling constant to about 300 Hz. This value is within the 250-730 Hz range indicative of four coordinate phosphorus.<sup>15</sup> As expected, the phenyl resonances are split by the phosphorus. As with other  $\text{R}_3\text{AlPR}_3$  adducts<sup>16,17</sup> no detectable splitting of the resonances for the alkyl groups bound to aluminum by phosphorus occurs.

The new organoaluminum phosphide  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$  has been prepared from appropriate reagents by the elimination of  $\text{SiMe}_4$  at  $160-180^\circ$  (eq. 3), hydrogen at  $70-80^\circ$  (eq. 4) and  $\text{KBr}$  (eq. 5) at room temperature. The best



synthetic reaction based on the ease of preparation and purification of reagents, ultimate purity and percent yield of purified product is given by eq. 3. The purified product  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$ , a colorless solid, is isolated in 78% yield, even though  $\text{SiMe}_4$  is formed in 95% yield. If reaction temperatures higher than  $160-180^\circ$ , i.e.  $180-200^\circ$ , are used for the preparative

reaction, reaction is considerably faster. However, products indicative of the decomposition of the trimethylsilylmethyl ligand,  $H_2$  and a solid product with a grey color are formed. The compound,<sup>7</sup>  $KAl(CH_2SiMe_3)_3H$ , also decomposes around  $200^\circ$ .

The potentially amphoteric ligand,  $(Me_3SiCH_2)_2AlPPh_2$ , exists as a mixture of monomeric and dimeric species in benzene solution according to cryoscopic molecular weight studies. The compounds  $Me_2AlPPh_2$  and  $Et_2AlPPh_2$  are dimers. All other organoaluminum phosphides exist as either dimers or trimers in solution.<sup>12</sup> The decreased extent of association for  $(Me_3SiCH_2)_2AlPPh_2$  is presumably due to the bulky  $CH_2SiMe_3$  ligand. The compound,  $(Me_3SiCH_2)_2AlPPh_2$  is the most soluble of the three organoaluminum phosphides discussed in this study, having moderate solubility in hexane and pentane and excellent solubility in aromatic hydrocarbons and diethyl ether. The spectral properties of these compounds are discussed later.

The compound  $(Me_3SiCH_2)_2AlPPh_2$  exhibits an unusual behavior upon heating. The purified product from the  $Al(CH_2SiMe_3)_3-PPh_2H$  reaction shows a solid to glass transformation at  $120-121^\circ$  and this glass melts at  $136-139^\circ$ . If this sample is then permitted to cool slowly to room temperature, crystals form which upon reheating melt at  $141-143^\circ$ , no glass transition is observed. The purified product from the  $Al(CH_2SiMe_3)_2H-PPh_2H$  reaction does not show the glass transition but melts sharply at  $146-147^\circ$ .

The compound  $Me_2AlPPh_2$  is also readily prepared by a variety of routes, in the presence or absence of a solvent (toluene) or excess alane. All of the reactions are quantitative. All products have the same  $^1H$  NMR and infrared spectral properties and analyses but they have slightly different melting points. The different melting points observed for the

products from the different reactions and/or conditions are probably related to the detailed nature and extent of association of the aluminum-phosphide rather than impurities. Some samples of product melt sharply at 228°, whereas others form a glass at temperatures around 200° and then melt at higher temperatures, within the range of 210 to 225°. Since these temperatures are not considered to reflect a material impurity, the melting point of this type of compound should not be used for product identification and purity. The details of the melting point behavior of different products are given in the Experimental Section. The compound  $\text{Et}_2\text{AlPPh}_2$  also exhibits this unusual glass transition at 123-126° prior to melting at 141-143°. Similar observations have been made for many other Group 3-5<sup>2,18,19</sup> and 3-6<sup>20</sup> compounds.

The aluminum-phosphorus dimers have been reported to react with Lewis bases to form adducts,  $\text{PPh}_2\text{AlR}_2 \cdot \text{Base}$ . The compound,  $\text{PPh}_2\text{AlEt}_2 \cdot \text{OEt}_2$ , was first described by Issleib and Deyling.<sup>21</sup> The compound was prepared by two different routes but the melting points of the two resulting solid products were slightly different, 148° and 151°. In a later paper Issleib and Krech<sup>22</sup> reported  $\text{PPh}_2\text{AlEt}_2 \cdot \text{OEt}_2$  as a solid with a melting point of 145-150°. This compound was characterized by analysis and curious cryoscopic molecular weight measurements in dioxane solution. Johnson, Larson and Dahl<sup>3</sup> also reported  $(\text{Et}_2\text{AlPPh}_2)_2$  (mp: 124-126°) and  $\text{PPh}_2\text{AlEt}_2 \cdot \text{OEt}_2$  (mp: 148°). The formulation of the etherate was supported by cryoscopic molecular weight measurements and hydrolyzable ethane data but analysis for P and Al were consistent with the formula  $\text{Et}_2\text{AlPPh}_2$ , not the etherate. We have attempted to repeat the synthesis of  $\text{PPh}_2\text{AlEt}_2 \cdot \text{OEt}_2$  but without success. All samples of  $\text{Et}_2\text{AlPPh}_2$  combined with  $\text{Et}_2\text{O}$  produce a solid which melts in the temperature range reported for  $\text{PPh}_2\text{AlEt}_2 \cdot \text{OEt}_2$ . However, mass balance experiments and

$^1\text{H}$  NMR spectra of products show conclusively that  $\text{Et}_2\text{O}$  does not react with  $\text{Et}_2\text{AlPPh}_2$  to form an adduct. Our results suggest that the earlier workers<sup>3,21,22</sup> were confused and misled by the unusual melting point behavior of  $\text{Et}_2\text{AlPPh}_2$ . The glass transition of  $\text{Et}_2\text{AlPPh}_2$  at  $123\text{--}126^\circ$  is the melting point previously reported<sup>3</sup> for  $\text{Et}_2\text{AlPPh}_2$ , whereas the melting point of  $\text{Et}_2\text{AlPPh}_2$  at  $141\text{--}143^\circ$  is the melting point reported for the proposed ether adduct.<sup>3,21,22</sup> Thus, neither  $\text{Et}_2\text{AlPPh}_2$  nor  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$  form stable diethyl ether adducts.


Our attempts to synthesize compounds<sup>23</sup> of the type  $\text{Cr}(\text{CO})_5(\text{PPh}_2\text{AlR}_2)$  from  $\text{KCr}(\text{CO})_5\text{PPh}_2 \cdot 2$  dioxane and  $\text{AlR}_2\text{Br}$  in tetrahydrofuran unexpectedly produced products containing a cleaved THF moiety,  $\text{Cr}(\text{CO})_5[\text{PPh}_2\text{C}_4\text{H}_8\text{OAlR}_2]$ . These results led us to question the claim that  $\text{PPh}_2\text{AlEt}_2 \cdot \text{THF}$  can be prepared from  $\text{LiPPh}_2$  and  $\text{AlEt}_2\text{Br}$  in THF. We have studied the reaction of  $\text{Et}_2\text{AlPPh}_2$  with excess THF at room temperature. If the excess THF is removed after 18 h, a sticky white solid is isolated. Its  $^1\text{H}$  NMR spectrum shows lines indicative of a cleaved THF moiety. Acid hydrolysis of the sticky white solid produces  $\text{Ph}_2\text{PC}_4\text{H}_8\text{OH}$  and a small amount of  $\text{PPh}_2\text{H}$ . The former compound has been previously obtained after hydrolysis of the product from  $\text{LiPPh}_2$  and THF.<sup>24</sup> The  $\text{PPh}_2\text{H}$  probably results from the hydrolysis of  $[\text{Et}_2\text{AlPPh}_2]_2$  which did not react with THF rather than  $\text{PPh}_2\text{AlEt}_2 \cdot \text{THF}$ . The compounds  $\text{Me}_2\text{AlPPh}_2$  and  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$  also cleave THF. The relative rates of cleavage of THF have been monitored by  $^1\text{H}$  NMR experiments and have been found to increase in the order  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2 > \text{Me}_2\text{AlPPh}_2 > \text{Et}_2\text{AlPPh}_2$ , with reaction ranging from one-half to essentially complete in 8 h. In all cases, the THF cleaved products  $\text{R}_2\text{AlOC}_4\text{H}_8\text{PPh}_2$  exhibit several alkylaluminum  $^1\text{H}$  NMR signals suggesting a mixture of variously associated oligomers.

Many organometallic main-group compounds<sup>25</sup> including various organo-aluminum compounds<sup>12</sup> and alkali metal phosphides,<sup>24,25</sup> cleave THF and other

ethers. In view of the generality of this reaction, it is surprising to note that there appears to be only one prior example involving a Group 3-5 compound as an ether cleaving reagent. The reaction mixture of  $B_2H_6$  and  $(CF_3)_2PH$  in dimethyl ether led to the formation of ether cleaved products at or below room temperature.<sup>27</sup> However, we believe that the THF cleavage reaction is probably more common in Group 3-5 chemistry than the literature suggests. The ether cleavage reaction appears to be very facile and the products are expected to have good thermodynamic stability.

The organoaluminum phosphides  $R_2AlPPh_2$  ( $R = Me, Et, CH_2SiMe_3$ ) are very reactive towards other common organic solvents as well as THF. Dichloromethane and chloroform solutions react to form grey precipitates after only 12 h at room temperature. Acetonitrile is reduced under equally mild conditions. Infrared spectra of the products from the reactions of all  $R_2AlPPh_2$  with  $CH_3CN$  show strong absorbances around  $1600\text{ cm}^{-1}$  attributable to the carbon-nitrogen double bond stretch. There are many examples of simple organoaluminum compounds such as  $AlMe_3$  and  $AlMe_2H$  reacting with  $CH_3CN$  to form dimeric aldimine derivatives  $[RCH = NAlMe_2]_2$ .<sup>28</sup> The reactions of  $Ar_2AlPPh_2$  ( $Ar = Ph, Me-C_6H_4$ ) with  $CH_3CN$  at or below room temperature have also been studied but adducts of the general formula  $Ar_2AlPPh_2 \cdot NCCH_3$  were observed.<sup>29</sup>

The  $^1H$  NMR spectra of  $[R_2AlPPh_2]_2$  ( $R = Me, Et, CH_2SiMe_3$ ) have been useful in revealing some details concerning the structures of these dimeric molecules in benzene solution. The data suggest that the four membered ring is nonplanar in solution with nonequivalent groups on aluminum and phosphorus.



The lines assigned to the hydrogens on the alkyl groups adjacent to aluminum are unexpectedly complex. The compound  $[\text{Me}_2\text{AlPPh}_2]_2$  in benzene solution exhibits an apparently symmetrical quartet at 90 or 100 MHz at  $\delta$  0.08 for the aluminum-methyl group. However, closer inspection of these lines gives a separation between the lines of 2.2, 2.1 and 2.2 Hz. At 60 MHz, the quartet is less symmetrical with separations of 2.1, 2.6 and 2.3 Hz. These observations suggest that the methyl groups are not equivalent and that the quartet is due to the superposition of two  $^{31}\text{P}$ - $^1\text{H}$  coupling patterns. Due to the fast relaxation of the  $^{27}\text{Al}$  nucleus, spin-spin coupling is not usually observed.<sup>30</sup> The alkyl region of the  $^1\text{H}$  NMR spectrum of  $[(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2]_2$  shows a singlet for the silicon methyl groups and an unsymmetrical quartet for the methylene groups at 90 MHz. Expansion reveals a shoulder on the side of the line at lowest field. At 60 MHz, the signal due to the aluminum methylene groups simplifies to a fairly symmetrical triplet with a coupling constant of 2.6 Hz. The spectrum is independent of concentration in the range 0.102-0.361 m. At these concentrations virtually all  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$  should be in the form of dimers, with nonequivalent aluminum alkyl groups. The alkyl region of the spectrum of  $[\text{Et}_2\text{AlPPh}_2]_2$  shows the expected splitting pattern for an ethyl group; a triplet at  $\delta$  1.31 for the methyl groups and a quartet at  $\delta$  0.80 for the methylene groups. The coupling constants are 7 Hz for both absorbances and are independent of spectrometer frequencies. However, further splitting of the methylene quartet is also observed. At all spectrometer frequencies, the methylene quartet lines are considerably broader than those due to the methyl groups. Expansion of the quartet does not reveal any details. The

phenyl resonances are not helpful in providing more information about any of the molecules. Two adjacent complex multiplets in the  $\delta$  7.7-7.1 region are observed. The complexity of these signals results from the overlap of the  $^1\text{H}$ - $^{31}\text{P}$  and  $^1\text{H}$ - $^1\text{H}$  coupling patterns for the ortho, meta and para ring hydrogens. The splitting of the lines for hydrogens on carbon adjacent to aluminum in the dimers may be explained by the existence of nonplanar, four-membered aluminum-phosphorus rings. A nonplanar dimer has been observed for the solid state structures of  $[\text{I}_2\text{BPPH}_2]_2$ <sup>31</sup> and  $[\text{Me}_2\text{AlN}(\text{iPr})\text{H}]_2$ .<sup>32</sup> The alkyl groups on aluminum are not equivalent and the hydrogens on the  $\alpha$ -carbon atoms would be split by two equivalent phosphorus atoms giving rise to a triplet for each of the two types of alkyl groups. Long range  $^{31}\text{P}$ - $^1\text{H}$  coupling is observed in other systems<sup>33</sup> but it has not been reported previously for other organoaluminum phosphides mainly because the NMR spectra have not been observed. The proposed structure is also consistent with the proton decoupled  $^{31}\text{P}$  NMR spectrum of  $[\text{Et}_2\text{AlPPh}_2]_2$ , which consist of a single absorption at  $\delta$  42.3 (downfield from  $\text{H}_3\text{PO}_4$ ) for two equivalent phosphorus atoms.

Acknowledgment. This work was supported in part by the Office of Naval Research.



References

1. Tessier-Youngs, C.; Bueno, C.; Beachley, O. T. Jr. Inorg. Chem. 1983, 22.
2. Coates, G. E.; Graham, J. J. Chem. Soc. 1963, 233.
3. Johnson, A. W.; Larson, W. D.; Dahl, G. H. Canad. J. Chem. 1966, 43, 1338.
4. Giurgiu, G.; Popescu, I.; Ciobame, A.; Bostan, M.; Voiculescu, N.; Roman, L. Rev. Roum. Chim. 1970, 15, 1581.
5. Beachley, O. T. Jr., Simmons, R. G. Inorg. Chem. 1980, 19, 1021.
6. Beachley, O. T. Jr.; Rusinko, R. N. Inorg. Chem. 1979, 18, 1966.
7. Beachley, O. T. Jr.; Tessier-Youngs, C.; Simmons, R. G.; Hallock, R. B. Inorg. Chem. 1982, 21, 1970.
8. Wartik, T.; Schlesinger, H. I. J. Amer. Chem. Soc. 1953, 75, 835.
9. a) Bianco, V. D.; Doronzo, S. Inorg. Synth. 1976, 16, 161.  
b) Kuchen, W.; Buchwald, H. Chem. Ber. 1958, 91, 2871.
10. Texas Alkyl Product Information.
11. Beg, M. A. A. Bull. Chem. Soc. Japan 1967, 40, 15.
12. Mole, T.; Jeffrey, E. A. "Organoaluminum Compounds." Elsevier, New York, 1972.
13. Al-Hashimi, S.; Smith, J. D. J. Organomet. Chem. 1978, 153, 253.
14. Adams, D. M. "Metal-Ligand and Related Vibrations," St. Martins Press, New York, 1968.
15. Brazier, J. F.; Houalla, D.; Koenig, M.; Wolf, R. Topics in Phos. Chem. 1976, 8, 99.
16. a) Clemens, D. F.; Sisler, H. H.; Brey, W. S. Jr. Inorg. Chem. 1966, 5, 527.  
b) Cullingworth, A. R.; Pidcock, A.; Smith, J. D. J. C. S. Chem. Comm. 1966, 89.  
c) Cohen, B. M.; Cullingworth, A. R.; Smith, J. D. J. Chem. Soc. (A) 1969, 2193.
17. Rosch, L.; Schmidh, W. Z. Anorg. Allg. Chem. 1976, 426, 99.
18. Beachley, O. T. Jr.; Coates, G. E.; Kohnstam, G. J. Chem. Soc. (A) 1965, 3248.

19. Beachley, O. T. Jr.; Coates, G. E. J. Chem. Soc. (A) 1965, 3241.
20. Turova, N. YA.; Kozunov, V.A.; Yanovskii, A. I.; Bokii, N. G.; Struchkov, Y. T.; Tarnopelskii, B. L. J. Inorg. Nucl. Chem. 1979, 41, 5 and references therein.
21. Issleib, K.; Deylig, H. J. Z. Naturforsch 1962, 17B, 198.
22. Issleib, K.; Krech, F. Z. Anorg. Allg. Chem. 1964, 328, 21.
23. Beachley, O. T. Jr.; Churchill, M. R.; Tessier-Youngs, C.; Youngs, W. unpublished observations.
24. Garner, A. Y.; Tedeschi, A. A. J. Amer. Chem. Soc. 1962, 84, 4734.
25. Coates, G. E.; Green, M. L. H.; Wade, K. "Organometallic Compounds", 3rd ed.; Methuen:London, 1967, Vol. 1 pp. 16, 75, 80-81.
26. Kosolapoff, G. M.; Maier, L. "Organic Phosphorous Compounds" Wiley-Interscience, New York, 1972, Vol. 1 pp. 42-44.
27. Burg, A. B.; Brendel, G. J. Amer. Chem. Soc. 1958, 80, 3198.
28. a) Wade, K.; Lloyd, J. E. J. Chem. Soc. 1965, 2662.  
b) Jennings, J. R.; Lloyd, J. E.; Wade, K. 1965, 5083.  
c) Wade, K.; Wyatt, B. K. J. Chem. Soc. (A) 1967, 1339.  
d) Wade, K.; Jennings, J. R. J. Chem. Soc. (A) 1967, 1333.
29. Giurgiu, D.; Popescu, L. Rev. Roum. Chim. 1976, 21, 1373.
30. Hinton, J. F.; Briggs, R. W. "NMR and the Periodic Table" (Harris, R. K.; Mann, B. E. editors) Academic Press. N.Y. 1978, p. 279.
31. a) Bullen, G. J.; Mallinson, P. R. J. Chem. Soc. Chem. Comm. 1969, 132.  
b) Mallison, K. B., Mann, F. G. J. Chem. Soc. 1964, 6121.
32. Amirkhalili, S.; Hitchcock, P. B.; Jenkins, A. D.; Nyathi, J. Z.; Smith, J. D. J. Chem. Soc. Dalton Trans. 1981, 377.
33. Nixon, J.; Pidcock, A. Ann. Report NMR Spec. 1969, 2, 345.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 North Quincy Street Arlington, Virginia 22217	2	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
U.S. Army Research Office Attn: CRD-AA-IP P. O. Box 12211 Research Triangle Park, N.C. 27709	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Mr. Vincent Schaper DTNSRDC Code 2803 Annapolis, Maryland 21402	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1

TECHNICAL REPORT DISTRIBUTION LIST, 053

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. M. F. Hawthorne Department of Chemistry University of California Los Angeles, California 90024	1	Dr. T. Marks Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. D. B. Brown Department of Chemistry University of Vermont Burlington, Vermont 05401	1	Dr. J. Zuckerman Department of Chemistry University of Oklahoma Norman, Oklahoma 73019	1
Dr. D. Venezky Chemistry Division Naval Research Laboratory Code 6130 Washington, D.C. 20375	1	<del>Professor O. T. Beachley</del> <del>Department of Chemistry</del> <del>State University of New York</del> <del>Buffalo, New York 14214</del>	1
Dr. John E. Jensen Hughes Research Laboratory 3011 Malibu Canyon Road Malibu, California 90265		Professor K. M. Nicholas Department of Chemistry Boston College Chestnut Hill, Massachusetts 02167	1
Dr. A. Cowley Department of Chemistry University of Texas Austin, Texas 78712	1	Professor R. Neilson Department of Chemistry Texas Christian University Fort Worth, Texas 76129	1
Dr. W. Hatfield Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1	Professor M. Newcomb Texas A&M University Department of Chemistry College Station, Texas 77843	1
Dr. M. H. Chisholm Department of Chemistry Indiana University Bloomington, Indiana 47401	1	Professor Richard Eisenberg Department of Chemistry University of Rochester Rochester, New York 14627	1
		Professor R. D. Archer University of Massachusetts Chemistry Department Amherst, Massachusetts 01003	

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 10	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Synthesis and Characterization of Organoaluminum Compounds Containing the Trimethylsilylmethyl Substituent, $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$ , $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$ and $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$ and a Reinvestigation of		5. TYPE OF REPORT & PERIOD COVERED Technical Report
7. AUTHOR(s) the Chemistry of $\text{Me}_2\text{AlPPh}_2$ and $\text{Et}_2\text{AlPPh}_2$ . O. T. Beachley, Jr. and Claire Tessier-Youngs		6. PERFORMING ORG. REPORT NUMBER N-0014-78C-0562
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-053-628
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, VA 22217		12. REPORT DATE February 8, 1983
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 22
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for Public Release, Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  Prepared for Publication in Inorganic Chemistry		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Organoaluminum compounds Organoaluminum hydride Organoaluminum bromide Organoaluminum phosphide Trimethylsilyl aluminum compounds		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The new amphoteric ligand, $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$ , has been prepared from the reactions of either $\text{Al}(\text{CH}_2\text{SiMe}_3)_3$ or $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$ with $\text{PPh}_2\text{H}$ or $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$ with $\text{KPh}_2$ and fully characterized by analysis, cryoscopic molecular weight measurements, IR and $^1\text{H}$ NMR data. This aluminum-phosphide is unique as it exists as a monomer-dimer equilibrium mixture in benzene solution. The syntheses and characterization of the new compounds $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{H}$ and $\text{Al}(\text{CH}_2\text{SiMe}_3)_2\text{Br}$ are also described. Since the chemistry of		

20.

of aluminum-phosphides as amphoteric ligands is of interest, the behavior of  $(\text{Me}_3\text{SiCH}_2)_2\text{AlPPh}_2$ ,  $(\text{Me}_2\text{AlPPh}_2)_2$  and  $(\text{Et}_2\text{AlPPh}_2)_2$  toward common solvents,  $\text{Et}_2\text{O}$ , THF and  $\text{CH}_3\text{CN}$ , was investigated. The aluminum phosphides readily cleave THF and reduce the triple bond of  $\text{CH}_3\text{CN}$  at room temperature. However, neither THF nor  $\text{Et}_2\text{O}$  form isolable adducts with aluminum phosphides. The unusual melting point behavior of aluminum phosphides is also discussed.